

A note on functions of exponential type

Antanas LAURINČIKAS* (VU, ŠU)
e-mail: antanas.laurincikas@maf.vu.lt

Let $0 < \theta_0 \leq \pi$. A function $f(s)$, $s = \sigma + it$, analytic in the closed angular region $|\arg s| \leq \theta_0$ is said to be of exponential type if

$$\limsup_{r \rightarrow \infty} \frac{\log |f(re^{i\theta})|}{r} < \infty$$

uniformly in θ , $|\theta| \leq \theta_0$.

Elements of the theory of functions of exponential type are applied in the investigation of the universality of Dirichlet series, see [1], Chapter 6. For example, the following result is very useful. Let \mathbb{C} be the complex plane, $\mathcal{B}(\mathbb{C})$ denote the class of Borel sets of the space \mathbb{C} , and let μ be a complex measure on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ with compact support contained in the strip $D = \{s \in \mathbb{C} : 1/2 < \sigma < 1\}$. Suppose that

$$\varrho(s) = \int_{\mathbb{C}} e^{-sz} d\mu(z)$$

and

$$\sum_p |\varrho(\log p)| < \infty.$$

Then, see Section 6.6.5 of [1],

$$\int_{\mathbb{C}} z^r d\mu(z) = 0, \quad r = 0, 1, 2, \dots$$

To study the universality, for example, of the function

$$\zeta^2(s)\zeta(2s-1),$$

where, as usual, $\zeta(s)$ denotes the Riemann zeta-function, we need an assertion similar to stated above for a complex measure μ_p depending on prime numbers p . More precisely, let μ be a

*Partially supported by Lithuanian State Studies and Science Foundation.

complex measure on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ with support contained in the strip $D_1 = \{s \in \mathbb{C} : 3/4 < \sigma < 1\}$ and let the function $h : \mathbb{C} \rightarrow \mathbb{C}$ be given by the formula

$$h(s) = \frac{s+1}{2}, \quad s \in \mathbb{C}.$$

Denote by

$$\mu_1(A) = \mu h^{-1}(A) = \mu(Ah^{-1}), \quad A \in \mathbb{C}.$$

Then we have that $\mu_p = 2\mu + a_p\mu_1$, $|a_p| = 1$, is a complex measure on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ with compact support contained in the strip D_1 . Define

$$\varrho_p(s) = \int_{\mathbb{C}} e^{-sz} d\mu_p(z).$$

Clearly, the function $\varrho_p(s)$ is a function of exponential type for each prime p .

Theorem 1. *Suppose that*

$$\sum_p |\varrho_p(\log p)| < \infty.$$

Then

$$\int_{\mathbb{C}} z^r d\mu(z) = 0, \quad r = 0, 1, 2, \dots$$

Proof. Define a number $\alpha > 0$ by

$$\sup_p \limsup_{t \rightarrow \infty} \frac{\log |\varrho_p(\pm it)|}{t} \leq \alpha,$$

and choose a number $\beta > 0$ satisfying the inequality $\alpha\beta < \pi$. Consider the set

$$A = \bigcup_{\substack{p \\ a_p \neq -1}} A_p,$$

where

$$A_p = \left\{ m \in \mathbb{N} : \exists r \in \left(\left(m - \frac{1}{4}\right)\beta, \left(m + \frac{1}{4}\right)\beta \right), \text{ and } |\varrho_p(r)| \leq e^{-r} \right\}.$$

Then we have

$$\sum_p |\varrho_p(\log p)| \geq \sum_{m \notin A} \sum'_m |\varrho_p(\log p)| \geq \sum_{m \notin A} \sum'_m \frac{1}{p}, \tag{1}$$

where \sum'_m denotes a sum over all prime numbers such that

$$\left(m - \frac{1}{4}\right)\beta < \log p \leq \left(m + \frac{1}{4}\right)\beta.$$

Using the well-known formula

$$\sum_{p \leq x} \frac{1}{p} = \log \log x + \gamma_0 + e^{-c_1 \sqrt{\log x}},$$

where γ_0 is Euler's constant and $c_1 > 0$, we find that

$$\begin{aligned} \sum'_m \frac{1}{p} &= \sum_{p \leq \exp\{m+1/4\}\beta} \frac{1}{p} - \sum_{p \leq \exp\{m-1/4\}\beta} \frac{1}{p} \\ &= \log \frac{m+1/4}{m-1/4} + B \exp\left\{-c_2 \left(\left(m - \frac{1}{4}\right)\beta\right)^{1/2}\right\} = \frac{1}{2m} + \frac{B}{m^2}. \end{aligned}$$

Here B is a number bounded by a constant. Hence and from (1) it follows that

$$\sum_{m \notin A} \left(\frac{1}{2m} + \frac{B}{m^2}\right) \leq \sum_p |\varrho_p(\log p)| < \infty,$$

and therefore

$$\sum_{m \notin A} \frac{1}{m} < \infty. \quad (2)$$

Let $A = \{a_1, a_2, \dots\}$ with $a_1 < a_2 < \dots$. Then (2) implies

$$\lim_{m \rightarrow \infty} \frac{a_m}{m} = 1. \quad (3)$$

By the definition of the set A there exist a sequence $\{\lambda_m\}$ and a prime number p_0 such that

$$\left(a_m - \frac{1}{4}\right)\beta < \lambda_m \leq \left(a_m + \frac{1}{4}\right)\beta$$

and

$$|\varrho_{p_0}(\lambda_m)| \leq e^{-\lambda_m}.$$

By (3)

$$\lim_{m \rightarrow \infty} \frac{\lambda_m}{m} = \beta,$$

and

$$\limsup_{m \rightarrow \infty} \frac{\log |\varrho_{p_0}(\lambda_m)|}{\lambda_m} \leq -1. \quad (4)$$

Now we will apply one version of the Bernstein theorem, Theorem 6.4.12 from [1]. Let $f(s)$ be an entire function of exponential type, and let $\{\lambda_m\}$ be a sequence of complex numbers. Let α, β and δ be positive real numbers such that

- a) $\limsup_{t \rightarrow \infty} \frac{\log |f(\pm it)|}{t} \leq \alpha;$
- b) $|\lambda_m - \lambda_n| \geq \delta |m - n|;$
- c) $\lim_{m \rightarrow \infty} \frac{\lambda_m}{m} = \beta;$
- d) $\alpha\beta < \pi.$

Then

$$\limsup_{m \rightarrow \infty} \frac{\log |f(\lambda_m)|}{|\lambda_m|} = \limsup_{r \rightarrow \infty} \frac{\log |f(r)|}{r}.$$

Consequently, by (4),

$$\limsup_{r \rightarrow \infty} \frac{\log |\varrho_{p_0}(r)|}{r} \leq -1. \quad (5)$$

Now we will use Lemma 6.4.10 from [1]. Let μ be a complex measure on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ with compact support contained in the half-plane $\sigma > \sigma_0$. Moreover, let

$$f(s) = \int_{\mathbb{C}} e^{sz} \, d\mu(z), \quad s \in \mathbb{C},$$

and $f(s) \not\equiv 0$. Then

$$\limsup_{r \rightarrow \infty} \frac{\log |f(r)|}{r} > \sigma_0.$$

Hence and from (5) we obtain that

$$\varrho_{p_0}(s) \equiv 0.$$

Differentiating the latter equation $r = 0, 1, 2, \dots$ times, and then putting $s = 0$, we find

$$\int_{\mathbb{C}} z^r \, d\mu_{p_0}(z) = 0. \quad (6)$$

Taking $r = 0$, we have

$$\int_{\mathbb{C}} d\mu_{p_0}(z) = 2 \int_{\mathbb{C}} d\mu(z) + a_{p_0} \int_{\mathbb{C}} d\mu_1(z) = (2 + a_{p_0}) \int_{\mathbb{C}} d\mu(z) = 0.$$

Thus, the equality

$$\int_{\mathbb{C}} z^r d\mu(z) = 0 \tag{7}$$

is true for $r = 0$. If $r = 1$, then (6) yields

$$\begin{aligned} 0 &= 2 \int_{\mathbb{C}} z d\mu(z) + a_{p_0} \int_{\mathbb{C}} z d\mu_1(z) \\ &= 2 \int_{\mathbb{C}} z d\mu(z) + a_{p_0} \int_{\mathbb{C}} (2z - 1) d\mu(z) \\ &= (2 + 2a_{p_0}) \int_{\mathbb{C}} z d\mu(z) - a_{p_0} \int_{\mathbb{C}} d\mu(z). \end{aligned}$$

Hence, in view of (7), we have that (7) is true for $r = 1$. The further proof is obtained in a standard induction way.

Theorem 1 is applied to obtain in a standard way the universality of the function $\zeta^2(s)\zeta(2s-1)$.

Theorem 2. *Let K be a compact subset of the strip D_1 with connected complement. Let $f(s)$ be a nonvanishing continuous function on K which is analytic in the interior of K . Then for every $\varepsilon > 0$*

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ \tau \in [0, T], \sup_{s \in K} |\zeta^2(s + i\tau)\zeta(2(s + i\tau) - 1) - f(s)| < \varepsilon \right\} > 0.$$

Here $\text{meas}\{A\}$ denotes the Lebesgue measure of the set A .

References

- [1] A. Laurinčikas, *Limit Theorems for the Riemann Zeta-Function*, Kluwer, Dordrecht (1996).

Pastaba apie eksponentinio tipo funkcijas

A. Laurinčikas

Straipsnyje nagrinėjama eksponentinio tipo funkcija, priklausanti nuo pirminių skaičių. Gauta tos funkcijos viena savybė, naudojama universalumo teorijoje.